

M.H.
09/463958
PRVPATENT- OCH REGISTRERINGSVERKET
Patentavdelningen

SOCO

REC'D 04 NOV 1999

WIPO

PCT

Intyg
Certificate

Härmed intygas att bifogade kopior överensstämmer med de handlingar som ursprungligen ingivits till Patent- och registreringsverket i nedannämnda ansökan.

This is to certify that the annexed is a true copy of the documents as originally filed with the Patent- and Registration Office in connection with the following patent application.

(71) Sökande AstaCarotene AB, Gustavsberg SE
Applicant (s)

(21) Patentansökningsnummer 9803541-3
Patent application number

(86) Ingivningsdatum 1998-10-16
Date of filing

Stockholm, 1999-09-29

För Patent- och registreringsverket
For the Patent- and Registration Office


Emma Johansson

Avgift
Fee 170:-

**PRIORITY
DOCUMENT**
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

**PATENT- OCH
REGISTRERINGSVERKET**
SWEDEN

Postadress/Adress
Box 5055
S-102 42 STOCKHOLM

Telefon/Phone
+46 8 782 25 00
Vx 08-782 25 00

Telex
17978
PATOREG S

Telefax
+46 8 666 02 86
08-666 02 86

29610/BN

TREATMENT OF DYSPEPSIA

5 The present invention relates to treatment of dyspepsia. The invention is particularly concerned with the use of at least one type of xanthophylls for the preparation of a medicament for prophylactic and/or therapeutic treatment of dyspepsia, and a method of treating dyspepsia. A preferred xanthophyll is astaxanthin.

10 Background of the invention

Dyspepsia, or indigestion, is one of the most common gastrointestinal disorders or diseases in both animals and humans. Dyspepsia is persistent or recurrent abdominal pain or abdominal discomfort centered in the upper abdomen. Dyspepsia refers to symptoms in the upper abdomen that are considered to be related to the upper alimentary tract. Often dyspepsia
15 is temporary and disappears spontaneously. In case alleviation of the symptoms is desired, ingestion of antacids is often the first choice of treatment. Antacids are normally not taken prophylactically.

However, there are several alternative therapeutic treatments of dyspepsia, and the present invention provides a new alternative, which may also be used for prophylactic
20 treatment.

Description of the invention

The present invention is directed to the use of at least one type of xanthophylls for the preparation of a medicament for prophylactic and/or therapeutic treatment of
25 dyspepsia.

In a preferred embodiment the xanthophyll is astaxanthin, especially astaxanthin in a form esterified with fatty acids. The astaxanthin in esterified form is suitably provided in the form of algal meal prepared from a culture of the alga *Haematococcus sp.*

The medicament may further comprise carbohydrate structures, such as
30 lipopolysaccharides, polysaccharides and glycoproteins, and/ or one or several different antioxidant(s), such as ascorbic acid (vitamin C) and tocopherol (vitamin E).

The medicament is preferably in the form of unit doses adapted for a daily dosage of xanthophyll(s) in the range of 0.05 to 1 mg per kg body weight of an individual, which is an animal or a human.

The invention is also directed to a method of prophylactic and/or therapeutic treatment of dyspepsia in an individual, which comprises administration to said individual of an dyspepsia-alleviating amount of a medicament comprising at least one type of xanthophylls.

5 In a preferred embodiment the xanthophyll is astaxanthin, especially astaxanthin in a form esterified with fatty acids. The astaxanthin in esterified form is suitably provided in the form of algal meal prepared from a culture of the alga *Haematococcus sp.*

The medicament may further comprise carbohydrate structures, such as lipopolysaccharides, polysaccharides and glycoproteins, and/ or one or several different
10 antioxidant(s), such as ascorbic acid (vitamin C) and tocopherol (vitamin E).

Preferably, the dyspepsia-alleviating amount of the medicament comprises xanthophyll(s) in the range of 0.05 to 1 mg per kg body weight of the individual, which is an animal or a human.

The at least one type of xanthophylls that are used in the present invention may
15 comprise a mixture of different types of xanthophylls or different forms of the same xanthophyll, such as a mixture of synthetic astaxanthin and naturally produced astaxanthin.

The mechanism of the prophylactic and therapeutic effect of the xanthophylls in the treatment of dyspepsia is not known, but it should be noted that they possess strong antioxidative properties and that they are soluble in fats and oils.

20 At present, the most preferred embodiment of the invention comprises algal meal having astaxanthin in esterified form with fatty acids dissolved in small droplets of naturally occurring oil and naturally occurring carbohydrate structures in the partially disrupted cell walls.

The medicament used in the invention may comprise additional ingredients
25 which are pharmacologically acceptable inactive or active in prophylactic and/or therapeutic use, such as flavoring agents.

The medicament may be presented in a separate unit dose or in mixture with food. Examples of separate unit doses are tablets, gelatin capsules and predetermined amounts of solutions, e. g. oil solutions, or emulsions, e.g. water-in- oil or oil-in-water emulsions.
30 Examples of foods in which the preparation of the invention may be incorporated is dairy products, such as yoghurt, chocolate and cereals.

Experiments

The experiments were conducted on 15 human volunteers experiencing symptoms of dyspepsia that did not disappear spontaneously. They were given 2 to 10 capsules per day containing algal meal produced by culturing of the algae *Haematococcus sp.* by AstaCarotene AB, Gustavsberg, Sweden. (These capsules are sold as an anti-oxidant and they contain 4 mg astaxanthin per capsule, with the instruction to take one capsule per day.)

This double to ten times the recommended dose, i.e. 8 - 40 mg astaxanthin per day, eliminated or drastically alleviated the symptoms of dyspepsia in all volunteers in 1 - 3 weeks, and no side effects were reported.

Astaxanthin from other sources, and other xanthophylls as well, are expected to be similarly useful for the purposes of the invention. An advantage of using astaxanthin from algae is, however, that the astaxanthin exists in a form esterified with fatty acids [Renström B. et al, 1981, Phytochem 20(11) :2561-2564], which esterified astaxanthin thereby is more stable during handling and storage than free astaxanthin.

15 -----

29610/BN

CLAIMS

1. Use of a at least one type of xanthophylls for the preparation of a medicament for prophylactic and/or therapeutic treatment of dyspepsia.
 - 5 2. Use according to claim 1, wherein the xanthophyll is astaxanthin.
 3. Use according to claim 2, wherein the astaxanthin is in a form esterified with fatty acids.
 4. Use according to claim 3, wherein the astaxanthin in esterified form is provided in the form of algal meal prepared from a culture of the alga *Haematococcus sp.*
 - 10 5. Use according to any one of claims 1 - 4, wherein the medicament further comprises carbohydrate structures.
 6. Use according to any one of claims 1 - 5, wherein the medicament further comprises (a) different antioxidant(s).
 7. Use according to any one of claims 1 - 6, wherein the medicament is in the form of unit doses adapted for a daily dosage of xanthophyll(s) in the range of 0.05 to 1 mg per kg body weight of an individual.
 - 15 8. Method of prophylactic and/or therapeutic treatment of dyspepsia in an individual, which comprises administration to said individual of an dyspepsia-alleviating amount of a medicament comprising at least one type of xanthophylls.
 - 20 9. Method according to claim 8, wherein the xanthophyll is astaxanthin.
 10. Method according to claim 9, wherein the astaxanthin is in a form esterified with fatty acids.
 11. Method according to claim 10, wherein the astaxanthin in esterified form is provided in the form of algal meal prepared from a culture of the alga *Haematococcus sp.*
 - 25 12. Method according claim 8, wherein the medicament further comprises carbohydrate structures.
 13. Method according to claim 8, wherein the medicament further comprises (a) different antioxidant(s).
 14. Method according to claim 8, wherein the dyspepsia-alleviating amount of the medicament comprises xanthophyll(s) in the range of 0.05 to 1 mg per kg body weight of the individual.
 - 30
-

29610/BN

ABSTRACT

Use of a at least one type of xanthophylls for the preparation of a medicament for prophylactic and/or therapeutic treatment of dyspepsia, is disclosed. The xanthophyll is preferably astaxanthin, e.g. astaxanthin is in a form esterified with fatty acids, such as in algal meal prepared from a culture of the alga *Haematococcus sp.* The medicament may further comprise carbohydrate structures and/or (a) different antioxidant(s). Further, a method of prophylactic and/or therapeutic treatment of dyspepsia in an individual, is described. The method comprises administration to said individual of an dyspepsia-alleviating amount of a medicament comprising at least one type of xanthophylls.
